arXiv:math/9310215v1 [math.FA] 11 Oct 1993

CONTRACTIVE PROJECTIONS IN NONATOMIC FUNCTION SPACES

BEATA RANDRIANANTOANINA

Abstract. We prove that there is no 1-complemented subspaces of finite codimension in separable rearrangament-invariant function spaces.

We study contractive projections onto finite-codimensional subspaces of real nonatomic function spaces. In general such projections are not common. It is well known that only in Hilbert space there exists a contractive projection onto every subspace of fixed finite codimension (cf. [1]).

The study of contractive projections (and more general projections with minimal norm) is important in approximation theory (cf. the survey of Cheney and Price [3]).

It is known that there is no 1-complemented subspaces of finite-codimension in $C[0,1]$ (Wulbert [13]) and in $L_1(\mu)$ if the measure μ is nonatomic ([6, Corollary IV.1.15], [4]). De Figueiredo and Karlovitz [5] (1970) proved that if μ is nonatomic then there is no 1-complemented hyperplanes in $L_p(\Omega, \mu)$, $1 < p < \infty$, $p \neq 2$, (cf. also [2]).

In this paper we prove that in rearrangement-invariant nonatomic function spaces not isometric to L_2 there is no 1-complemented subspaces of any finite codimension.

We use the terminology and notation as in [9].

Our method of proof is surprisingly simple – it is based on the following observation:

Proposition 1. (*cf.* [7, 11]) *In a real Banach space* X *if* P *is a projection then* $\|I - P\| = 1$ *(where* I *denotes identity operator) if and only if* $x^*(Px) \geq 0$ *for all* $x \in X$ *and* $x^* \in X^*$ *norming for* x.

In [7, Theorem 4.3] (cf. [8, 10]) Kalton and the author proved the nonexistence of 1-complemented hyperplanes in a wide class of nonatomic function spaces. However the original theorem uses special technical frazeology so we state it below in the language of projections:

1991 *Mathematics Subject Classification.* 46B,46E.

Theorem 2. *Suppose X is a real order-continuous Köthe function space on (Ω, μ) and μ is nonatomic. Then the hyperplane H in X is 1-complemented if and only if there exists a nonnegative measurable function w with* $\operatorname{supp} w = B = \operatorname{supp} f$, *where $f \in H^{\perp} \subset X^*$, so that for any $x \in X$ with* $\operatorname{supp} x \subset B$

$$\|x\| = (\int |x|^2 w \, d\mu)^{1/2}.$$

Hence there is no 1-complemented hyperplanes in X unless L_2 is isometric to a band in X. In particular there is no 1-complemented hyperplanes in separable r.i. spaces on $[0, 1]$ ([7, Theorem 4.4]).

As a part of proof of Theorem 2 we proved the following fact which we state separately for the future use.

Proposition 3. ([10, Proposition 2.8]) *Let X be an order-continuous Köthe function space on (Ω, μ), where μ is nonatomic. Suppose that the set*

$$\Lambda = \left\{ \frac{x^*}{x} \ : x \in X, x^* \in X^* \textit{ norming for } x \right\}$$

is one-dimensional i.e. $\Lambda \subset \{aw : a \in \mathbb{R}\}$ for some $w \in L_0(\Omega, \mu)$. Then X is isometric to $L_2(w d\mu)$.

Now we are ready to prove our main result.

Theorem 4. *Suppose μ is nonatomic and X is a separable r.i. space on $([0, 1], \mu)$ not isometric to L_2. Then there is no 1-complemented subspaces of any finite codimension in X.*

For the proof we need the following measure-theoretic lemma.

Lemma 5. *Suppose μ is nonatomic and suppose $f_1, \dots, f_n, g_1, \dots, g_n \in L_1(\mu)$ are such that $g_1, \dots, g_n$ are linearly independent and*

$$\sum_{i=1}^{n} \left(\int h f_j \, d\mu \right) \left(\int h g_j \, d\mu \right) \geq 0$$

whenever $|h| = 1$ a.e. Then $\{f_j\}_{j=1}^n \subset \operatorname{span} \{g_j\}_{j=1}^n$.

Proof. Consider the subset Γ of $\mathbb{R}^{2n}$ of all $2n$-tuples $(a_1, b_1, \dots, a_n, b_n)$ such that for some $h \in L_\infty(\mu)$ with $|h| = 1$ a.e. we have $a_j = \int hf_j \, d\mu$, $b_j = \int hg_j \, d\mu$, $j = 1, \dots, n$. Let $H = \operatorname{span} \Gamma \subset \mathbb{R}^{2n}$.

We immediately see that $\Gamma = -\Gamma$ and that by Liapunoff's theorem ([12]) Γ is convex. Hence for every $(s_1, t_1, \dots, s_n, t_n) \in H$ we have $\sum_{j=1}^{n} s_j t_j \geq 0$.

On the other hand consider $V = \{v_k = e_{2k-1} - e_{2k} : k = 1, \dots, n\} \subset \mathbb{R}^{2n}$ where e_j denotes the natural basis of $\mathbb{R}^{2n}$. Clearly $V \cap H = \{0\}$ so $\dim H \leq n$. Therefore there exist $c_1, \dots, c_n \in \mathbb{R}$ such that $s_i = \sum_{j=1}^{n} c_j t_j$ for $i = 1, \dots, n$, for every $(s_1, t_1, \dots, s_n, t_n) \in H$. In particular $\int hf_i \, d\mu = \int h \left(\sum_{j=1}^{n} c_j g_j \right) d\mu$ for all h with $|h| = 1$ a.e. and the lemma follows. □

Proof of Theorem 4. Suppose that F is a closed linear subspace of codimension n in X and let $u_1, \dots, u_n$ be linearly independent functions in X such that $\operatorname{span}\{F, u_1, \dots, u_n\} = X$. Denote $P : X \longrightarrow F$ a contractive projection onto F and consider $Q = I - P$. Then $Q = \sum_{j=1}^{n} f_j \otimes u_j$ for some linearly independent $f_1, \dots, f_n \in X^*$.

By Proposition 1 for any $x \in X$ and $x^* \in X^*$ norming for x $x^*(Qx) \geq 0$. Next for any h with $|h| = 1$ a.e. hx^* is norming for hx if x^* is norming for x. Hence $hx^*(Q(hx)) \geq 0$ i.e.

$$\sum_{j=1}^{n} \left(\int f_j hx \, d\mu \right) \left(\int u_j hx^* \, d\mu \right) \geq 0.$$

By Lemma 5 $f_1 x \in \operatorname{span}\{u_j x^* : j = 1, \dots, n\}$. So if $B = \operatorname{supp} f_1$ ($\mu(B) > 0$) then $\left.\frac{x}{x^*}\right|_B \in \operatorname{span}\left\{ \frac{u_j}{f_1} : j = 1, \dots, n \right\}$. By re-arrangement invariance of X for every measure preserving map $\sigma : [0,1] \longrightarrow [0,1]$ $x^* \circ \sigma$ is norming for $x \circ \sigma$ and so

$$\left.\frac{x \circ \sigma}{x^* \circ \sigma}\right|_B \in \operatorname{span}\left\{ \frac{u_j}{f_1} : j = 1, \dots, n \right\}$$

i.e. the set $\left\{ \left.\left[\left(\frac{x}{x^*} \right) \circ \sigma \right]\right|_B \mid \sigma : [0,1] \longrightarrow [0,1] \text{ measure preserving} \right\}$ is finite dimensional which is impossible unless $\frac{x}{x^*}$ is a constant. But then by Proposition 3 we conclude that X is isometric to $L_2[0,1]$ contrary to our assumption. □

Remark . Notice that in the proof of Theorem 4 we use re-arrangement invariance of X only in the final step to conclude that if the set $\left\{ \frac{x^*}{x} : x \in X, x^* \in X^* \text{ norming for } x \right\}$ is finite-dimensional then it is one-dimensional. Similar conclusion is true also in spaces of the form $X(Y)$, $X_1(X_2(\ldots(X_m)\ldots))$, where X, Y, X_j are r.i. and Theorem 4 holds as stated also for those spaces.

Acknowledgments . I wish to express my gratitude to Professor Nigel Kalton for his interest in this work and many valuable disscussions.

References

[1] D. Amir. *Characterizations of inner product spaces.* Birkhäuser Verlag, Basel, Boston Stuttgart, 1986.

[2] B. Beauzamy and B. Maurey. *Points minimaux et ensembles optimaux dans les espaces de Banach.* J. Functional Anal. **24** (1977) 107–139.

[3] E. W. Cheney and K. H. Price. *Minimal projections.* In: A. Talbot, editor, *Approximation Theory, Proceedings of a symposium held at Lancaster, July 1969*, pages 261 – 289. Academic Press, 1970.

[4] R. G. Douglas. *Contractive projections on an L_1 space.* Pacific J. Math. **15** (1965) 443–462.

[5] D.G. de Figueiredo and L. A. Karlovitz. *On the extention of contractions of normed spaces.* In: *Nonlinear Functional Analysis Proceedings of symposia in Pure Mathematics, Volume 18, Part 1*, pages 95–104, Providence, 1970. Amer. Math. Soc.

[6] P. Harmand, D. Werner, and W. Werner. *M-ideals in Banach Spaces and Banach Algebras.* Lecture Notes in Math. 1547. Springer, Berlin-Heidelberg-New York, 1993.

[7] N. J. Kalton and B. Randrianantoanina. *Surjective isometries of rearrangement-invariant spaces.* to appear in Quart. J. Math. Oxford.

[8] N. J. Kalton and B. Randrianantoanina. *Isometries of rearrangement-invariant spaces.* C. R. Acad. Sc. Paris **316** (1993) 351–355.

[9] J. Lindenstraus and L. Tzafriri. *Classical Banach spaces, Vol. 2, Function spaces.* Springer–Verlag, Berlin–Heidelberg–New York, 1979.

[10] B. Randrianantoanina. *Isometries of function spaces, Ph.D. Dissertation.* University of Missouri–Columbia, 1993.

[11] H. P. Rosenthal. *Contractively complemented subspaces of Banach spaces with reverse monotone (transfinite) bases.* Longhorn Notes, The University of Texas Functional Analysis Seminar (1984-5) 1–14.

[12] W. Rudin. *Functional Analysis.* McGraw–Hill, New York, second edition, 1991.

[13] D. E. Wulbert. *Projections of norm 1 on $C(X)$.* Notices Amer. Math. Soc. **15** (1968) 362.

Department of Mathematics, Bowling Green State University, Bowling Green, OH 43403

E-mail address: `brandri@andy.bgsu.edu`